

# Scientists' warning: six key points where biodiversity can improve climate change mitigation

Cássio Cardoso Pereira , Walisson Kenedy-Siqueira , Daniel Negreiros , Stephannie Fernandes , Milton Barbosa , Fernando Figueiredo Goulart , Simone Athayde , Christopher Wolf , Ian J. Harrison , Matthew G. Betts , Jennifer S. Powers , Rodolfo Dirzo , William J. Ripple , Philip M. Fearnside  and Geraldo Wilson Fernandes 

Cássio Cardoso Pereira ([cassiocardosopereira@gmail.com](mailto:cassiocardosopereira@gmail.com)), Walisson Kenedy-Siqueira, Daniel Negreiros, Fernando Figueiredo Goulart, and Geraldo Wilson Fernandes are affiliated with the Knowledge Center for Biodiversity, in Belo Horizonte, Minas Gerais, in Brazil and with the Ecologia Evolutiva & Biodiversidade, Departamento de Genética, Ecologia e Evolução, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil. Stephannie Fernandes and Simone Athayde are affiliated with the Department of Global and Sociocultural Studies and with the Kimberly Green Latin American and Caribbean Center, at Florida International University, in Miami, Florida, in the United States. Milton Barbosa is affiliated with the Environmental Change Institute, in the School of Geography and the Environment, at the University of Oxford, in Oxford, England, in the United Kingdom. Christopher Wolf, Matthew G. Betts, and William J. Ripple are affiliated with the Department of Forest Ecosystems and Society at Oregon State University, in Corvallis, Oregon, in the United States. Ian J. Harrison is affiliated with the Moore Center for Science, at Conservation International, and with the Freshwater Conservation Committee, IUCN Species Survival Commission, in Arlington, Virginia, in the United States. Jennifer S. Powers is affiliated with the Department of Plant and Microbial Biology at the University of Minnesota, in St. Paul, Minnesota, in the United States. Rodolfo Dirzo is affiliated with the Department of Biology and Earth Systems Science at Stanford University, in Stanford, California, in the United States. Philip M. Fearnside is affiliated with the Instituto Nacional de Pesquisas da Amazônia, in Manaus, Amazonas, in Brazil.

The focus on removing carbon dioxide from the atmosphere reflects increased public attention to climate change that potentially comes at the expense of other biodiversity challenges (Pereira et al. 2023a). This asymmetry between environmental agendas harms not only biodiversity but also climate-change mitigation because environmental issues are inexorably intertwined (Pörtner et al. 2023).

Climate-change-related extreme weather events and disasters are emerging across the planet, resulting in unprecedented economic, social, and ecological losses (Ripple et al. 2017). Solving the climate crisis is urgent, but the net-zero carbon emission commitments for 2050 are likely to fail if biodiversity issues are not fully integrated into the international climate agenda. It is well understood that biodiversity promotes multiple socioenvironmental services and benefits, including water and air quality, crop pollination, food security, human health and well-being, and protection from soil erosion. Climate change can accelerate biodiversity loss, and the associated ecosystem degradation undermines ecosystem resilience and reduces climate-change mitigation by reducing carbon sequestration (Pörtner et al. 2023). This exacerbates the impact of extreme weather events, resulting in increased vulnerability and socioeconomic losses.

Given these links, there is increasing recognition of the need for a more integrated approach to tackle the climate and biodiversity crises. Below, we list six ways in which the protection, conservation, and restoration of biodiversity can improve climate-change mitigation.

## 1. Conservation of carbon stocks and sinks

The current approaches are unlikely to deliver the climate benefits they promise if native ecosystems are replaced by exotic monospecific stands and if biodiversity and ecosystem functioning are not part of planning. The misguided replacement of native vegetation with tree plantations as carbon sinks results in the loss

of tropical forests, savannas, and grasslands to exotic stands of *Pinus* or *Eucalyptus*. This is a serious mistake; each ecosystem has its own importance and must be preserved as it is, especially because much of the carbon is stored in the soil rather than in trees. For example, the preserved soil of a grassland acts as a carbon sink, but when vegetation is removed or replaced by a monospecific plantation, the sink can become a source. We must expand the protection of natural ecosystems in order to promote the maintenance of carbon stocks (figure 1).

Protecting carbon stocks in ecosystems must be the first priority both for carbon stocks and for biodiversity (Pörtner et al. 2023). Native forests in the Amazon, the Congo Basin, and Southeast Asia are particularly important because of the co-occurrence of high carbon and biodiversity. Priorities vary among locations and at each moment in history at any given region. Because financial and human resources are always limited, these resources must be used to stem further losses before the restoration of degraded lands becomes a priority, since the carbon and biodiversity benefits of avoiding deforestation are very much greater than those of planting trees both per hectare and per dollar invested. This is the case when resources from the national budget or from international sources are allocated in countries where substantial areas of native vegetation still exist and are rapidly being lost, as in Brazil. However, in countries or in subnational units (such as the Brazilian states of São Paulo and Minas Gerais) where unprotected native vegetation is relatively scarce and where the governments of these countries or states have their own financial resources, the restoration of ecosystems will become a priority for environmental funds.

## 2. Biodiverse restoration

Where restoration is the appropriate priority, the way that it is done has important consequences. Many countries have committed to restoring degraded land, with promises totaling

Received: January 29, 2024. Revised: March 20, 2024. Accepted: April 5, 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of the American Institute of Biological Sciences.

All rights reserved. For permissions, please e-mail: [journals.permissions@oup.com](mailto:journals.permissions@oup.com)



**Figure 1.** Six key points in which tackling the biodiversity crisis can contribute to an effective solution to the climate emergency (top). Below it is shown how these six key points can be translated into targets for solving the twin crises together. Illustration: Walisson Kenedy-Siqueira®.

millions of hectares by 2030. But restoration takes much more than planting trees and covering bare land with any type of vegetation. Restoration projects have generally used a standard for all types of ecosystems, with little species diversity and without even knowing the vegetation neighboring the location where the restoration takes place (Toma et al. 2024). We are creating new ecosystems that fail to meet one of the most important objectives

of restoration: increasing environmental connectivity. By introducing a limited number of nonnative species into a given region, we can inadvertently reduce the ecological functionality of the environment, making it more homogeneous and less diverse. Only restoration with a diverse array of native species can more quickly promote environmental connectivity and restore the benefits that ecosystems can provide to humans (figure 1; Toma et al. 2024).

### 3. Integrated conservation of local fauna and flora

The conservation of wild animals and their roles in ecosystems are key components of natural solutions to climate change, which can enhance the ability to prevent climate warming beyond 1.5 degrees Celsius, potentially sequestering 6.5 petagrams of carbon per year (Schmitz et al. 2023). This may occur because of the role of animals in helping to mitigate and adapt to climate change and causes alteration in the fire regime and in the microclimate, as well as increasing carbon stocks, trophic complexity, the heterogeneity of habitats, and the pollination, dispersal, and propagation of plants (Pörtner et al. 2023). Conservation of forests must be integrated with the conservation of their resident fauna so that we can achieve our global climate mitigation goals (figure 1).

### 4. Use only existing areas of agriculture, pasture, and silviculture

Despite the importance of various kinds of plantations for human livelihoods, we must recognize that the expansion of these plantations is among the main drivers of fragmentation, loss of biodiversity and habitat, soil degradation, and impacts on nonclimate ecosystem services, such as water supply. Forest loss is disproportionately affecting biodiversity in landscapes around the world (Betts et al. 2017). Improved land management in existing areas of agriculture, ranching, and silviculture could sequester an additional 13.7 petagrams of carbon per year (Sha et al. 2022). The world's existing areas of plantations are sufficient for the human population to subsist, and there is no need to devastate new natural areas for cultivation. However, a lack of food can result from poor land-use choices, from economic inequalities impeding food purchase, and from transport barriers imposed by wars and natural disasters. We urge policymakers not to expand existing planted areas in order to slow the loss of terrestrial and aquatic ecosystems around the world (figure 1).

### 5. Incorporate biodiversity into business models

Solutions to the joint climate–biodiversity crises may partly lie in the private sector. Decades of experience have helped governments and corporations understand how to incorporate climate change into their business models, but economic incentives for sociobiodiversity conservation lag far behind. A high-level analysis by Fortune Global 500 shows that 83% of companies have climate-related goals, particularly in the transportation sector, whereas only 51% of companies recognize biodiversity loss in some way, and only 5% have set quantified targets beyond mere recognition (Claes et al. 2022). Businesses and financial institutions need to define sustainability more precisely in terms of biodiversity conservation, and incentives must be provided to do so. The Global Environment Facility (GEF) is on the right track by allocating essential resources to the Convention on Biological Diversity (CBD) (Pereira et al. 2023b). Likewise, the private sector can achieve a Net Positive Impact (NPI), which, over a quantified timescale, can outweigh the biodiversity disturbances and damage associated with the NPI activities (figure 1).

### 6. Joint biodiversity–climate conferences of the parties

To achieve net-zero emissions, it is necessary to align policies and actions across sectors and scales (Pettorelli et al. 2021). In 2021 the first joint report produced by the Intergovernmental Panel on Climate Change (IPCC) and the Intergovernmental Science–Policy Platform on Biodiversity and Ecosystem Services (IPBES) concluded that the world must tackle climate change and biodiversity loss together if either issue is to be successfully addressed. In this sense, the integration of environmental conferences would increase synergies between multilateral environmental agreements and international institutions. This would foster collaboration between experts on related topics, aligning methods and models and leading to a better assessment of trade-offs and interactions between different types of environmental impacts and policies (figure 1).

### Conclusions

To save the planet, conservation and the restoration of ecosystems must be considered to keep global warming below 1.5 degrees Celsius and to ensure a livable future. To achieve a sustainable future, we urgently need to commit to the key points presented here. Protecting a livable future will require rapid commitment not only from countries through actions in their national territories but also from emerging coalitions and governance models at all levels. Finally, we call on the media to foster a more balanced communication strategy to draw society's general attention to the role of biodiversity in addressing the climate-change crisis.

### Acknowledgments

We thank Jos Barlow for his uplifting comments that helped us with our message. CCP thanks scholarship from CNPq (grant no. 173800/2023–8). WKS thanks Project Peld-CRSC (CNPq-MCTI) for grant support. DN thanks scholarship from CNPq (grant no. 151341/2023–0). FFG thanks FINEP for the DTI (Development, Technology and Innovation) scholarship. PMF thanks Fundação de Amparo à Pesquisa do Estado de São Paulo (grant no. 2020/08916–8), Fundação de Amparo à Pesquisa do Estado do Amazonas (grant no. 0102016301000289/2021–33), FINEP/Rede CLIMA (grant no. 01.13.0353–00), and Conselho Nacional do Desenvolvimento Científico e Tecnológico (CNPq; grants no. 312450/2021–4 and no. 406941/2022–0). GWF thanks Knowledge Center for Biodiversity (CNPq; grant no. 406757/20224), CNPq-MCTI, and FAPEMIG for grant support.

### References cited

- Betts MG, Wolf C, Ripple WJ, Phalan B, Millers KA, Duarte A, Butchart SHM, Levi T. 2017. Global forest loss disproportionately erodes biodiversity in intact landscapes. *Nature* 547: 441–444.
- Claes J, Erben I, Hopman D, Jayaram K, Katz J, Aken TV. 2022. Where the world's largest companies stand on nature. *McKinsey Sustainability* (13 September 2022). [www.mckinsey.com/capabilities/sustainability/our-insights/where-the-worlds-largest-companies-stand-on-nature](https://www.mckinsey.com/capabilities/sustainability/our-insights/where-the-worlds-largest-companies-stand-on-nature).
- Pereira CC, Negreiros D, Barbosa M, Goulart FF, Dias RDL, Melillo MC, Camarota F, Pimenta MA, Cruz M, Fernandes GW. 2023a. Has climate change hijacked the environmental agenda? *Nature Conservation* 53: 157–164.

- Pereira CC, Fernandes GW, Negreiros D, Kenedy-Siqueira W, Fernandes S, Fearnside PM. 2023b. Hope for funding biodiversity efforts. *Science* 382: 383–384.
- Pettorelli N, Graham NAJ, Seddon N, Bustamante MC, Lowton MJ, Sutherland WJ, Koldewey HJ, Prentice HC, Barlow J. 2021. Time to integrate global climate change and biodiversity science-policy agendas. *Journal of Applied Ecology* 58: 2384–2393.
- Pörtner H-O, et al. 2023. Overcoming the coupled climate and biodiversity crises and their societal impacts. *Science* 380: eabl4881.
- Ripple WJ, et al. 2017. World scientists' warning to humanity: A second notice. *BioScience* 67: 1026–1028.
- Schmitz OJ, et al. 2023. Trophic rewilding can expand natural climate solutions. *Nature Climate Change* 13: 324–333.
- Sha Z, Bai Y, Li R, Lan H, Zhang X, Li J, Liu X, Chang S, Xie Y. 2022. The global carbon sink potential of terrestrial vegetation can be increased substantially by optimal land management. *Communications Earth and Environment* 3: 8.
- Toma TSP, et al. 2024. Aim for heterogeneous biodiversity restoration. *Science* 383: 376–376.